



NATIONAL UNIVERSITY OF SCIENCE AND TECHNOLOGY

FACULTY OF APPLIED SCIENCE

DEPARTMENT OF APPLIED CHEMISTRY

TRANSPORT PHENOMENA

SCH 2108

Supplementary Examination Paper

August 2015

This examination paper consists of 4 pages

Time Allowed: 3 hours

Total Marks: 100

Special Requirements: None

Examiner's Name: Mr. B. Nyoni

INSTRUCTIONS

1. Answer all questions in Section A and any other three questions from Section B
2. Each question carries 20 marks
3. Show steps clearly in any calculation
4. Start the answers for each question on a fresh page
5. Use of calculators is permissible

MARK ALLOCATION

QUESTION	MARKS
1.	20
2.	20
3.	20
4.	20
5.	20
TOTAL	100

Copyright: National University of Science and Technology, 2014

SECTION A

1 (a) Define the term '*dimension*' and list the seven basic quantities and their SI units. [10]

(b) What are the physical meanings of the following dimensionless groups:

(i) Reynolds number

(ii) Prandtl number

(iii) Nusselt number [10]

2 (a) What do you understand by the term '*a dimensionally consistent equation*'. [5]

(b) Check the dimensional consistency of the following empirical equation for a heat-transfer coefficient.

$$h_i = 0.023G^{0.8}k^{0.67}c_p^{0.33}D^{-0.2}\mu^{-0.47}$$

given h_i = heat transfer coefficient ($\text{W}/\text{m}^2\cdot^\circ\text{C}$)

G = mass velocity ($\text{kg}/\text{s}\cdot\text{m}^2$)

k = thermal conductivity ($\text{W}/\text{m}\cdot^\circ\text{C}$)

c_p = specific heat ($\text{J}/\text{g}\cdot^\circ\text{C}$)

D = diameter

μ = absolute viscosity ($\text{kg}/\text{m}\cdot\text{s}$) [12]

(c) What is the heat transfer coefficient, given the following data:

$$G = 54 \text{ kg}/\text{s}\cdot\text{m}^2$$

$$k = 0.12 \text{ W}/\text{m}\cdot^\circ\text{C}$$

$$c_p = 4.2 \text{ J}/\text{g}\cdot^\circ\text{C}$$

$$D = 0.11 \text{ m}$$

$$\mu = 0.034 \text{ kg}/\text{m}\cdot\text{s} [3]$$

SECTION B

3 (a) Define the following terms;

i) Viscosity

ii) Density

[4]

(b) Describe and explain the difference between a Newtonian and non-Newtonian fluid and give an example of each. [6]


(c) Prove that the resistance (F) of a sphere with diameter (D) moving at a constant speed (u) through a fluid of density (ρ) and viscosity (μ) is given by

$$F = \left(\frac{\mu}{\rho^2} \right) f \left(\frac{\rho u D}{\mu} \right) \quad [10]$$

4 (a) Derive the basic equation of fluid statics [8]

(b) Crude oil, specific gravity = 0.887, flows through the piping shown below, where the diameters are, pipe A is 50 mm, pipe B is 75 mm, and each of pipes C is 38 mm. An equal quantity of liquid flows through each of the pipes C. The flow through pipe A is 6.65 m³/h. Calculate (a) the mass flowrate through each pipe, (b) the average linear velocity through each pipe.

[Take $\pi = 3.142$]



[12]

- 5 (a) Discuss the mechanisms of heat transfer by which the sun transfers its energy to the earth's surface. [6]
- (b) Describe any two heat exchanger types of your choice. [4]
- (c) Calculate the rate of heat transfer by solar radiation on a flat concrete roof of a building, 8 m by 9 m, if the surface temperature of the roof is 330 K. The emissivity of concrete at 330 K is 0.89, whilst the total absorptivity of solar radiation (sun temperature = 5500 K) at this temperature is 0.60. [10]
- 6 (a) State the following;
- (i) Fourier's law of heat conduction
 - (ii) Fick's law of diffusion [6]
- (b) A tube or bridge of a gel solution of 1.05 wt% agar in water is at 278K and 0.04m long. It connects two agitated solutions of urea in water. The urea concentration in the first solution is 0.2 mol/litre and is 0 in the other. $D_{AB} = 0.727 \times 10^{-9} \text{ m}^2/\text{s}$.
- (i) Draw a diagram of the described setup
 - (ii) Calculate the flux of the urea in $\text{kmol}/\text{m}^2\text{s}$ at steady state. [14]

End of Question Paper!!!