

FACULTY OF APPLIED SCIENCES
DEPARTMENT OF APPLIED MATHEMATICS
SMA 2106: PROBABILITY THEORY

AUGUST 2004 SUPPLEMENTARY

Time : 3 hours

Candidates should attempt ALL questions from Section A and ANY THREE questions from Section B

SECTION A: [25 marks]

Answer ALL questions from this section.

A1. Let (Ω, \mathcal{F}, P) be a probability space and A be a subset of Ω . The collection \mathcal{G} of all subsets of Ω of the form $G = A \cap F$, $F \in \mathcal{F}$, is called the trace of \mathcal{F} on A .

(a) Show that the trace of \mathcal{F} on A is a σ -algebra in A [6]

(b) If $A \in \mathcal{F}$ and $P(A) > 0$, define a set function Q_A on \mathcal{G} by $Q_A(G) = \frac{P(G)}{P(A)}$. [6]

Show that Q_A is a probability measure on \mathcal{G} .

A2. Find the value of k for which the following function is a joint density function of X and Y :

$$f_{X,Y}(x,y) = \begin{cases} k(6-x-y) & \text{if } (x,y) \in \{(x,y) \in \mathbb{R}^2 : x > 0, x+y \leq 6, x+2y \geq 4\} \\ 0 & \text{otherwise} \end{cases} \quad [6]$$

A3. An experiment consists of throwing two unbiased die and let S be the score, i.e the sum of the numbers that appear on both die and let π be the product of them. Define the following random variables X and Y as follows:

$$X = \begin{cases} 1 & \text{if } S \text{ is even} \\ 0 & \text{if } S \text{ is odd} \end{cases}$$

$$Y = \begin{cases} 1 & \text{if } \pi \text{ is even} \\ 0 & \text{if } \pi \text{ is odd} \end{cases}$$

- (a) Describe the distribution of the random vector (X, Y) . [4]
 (b) Calculate the distribution function $F_{X,Y}(x, y)$. [3]

SECTION B: [75 marks]

Answer **ANY THREE** questions from this section. Each question carries 25 marks.

- B4.** (a) The joint probability density function of (X, Y) is given by

$$f_{X,Y}(x, y) = \begin{cases} c(xy + y^2) & \text{if } 0 \leq x \leq 2; 0 \leq y \leq 2 \\ 0 & \text{otherwise} \end{cases}$$

- (i) Find c [2]
 (ii) Calculate $P(X + Y < 2)$ [5]
 (iii) Calculate the coefficient of correlation ρ_{XY} . [6]
- (b) Two points A and B are randomly selected on the circumference of a circle, centre origin O and radius r . Calculate the expectation of the area of triangle AOB. [8]
- (c) A random vector (X, Y) has the following characteristics:
 $E[X] = -1$; $E[Y] = 1$; $\text{Var}[X] = 4$ and $\text{Var}[Y] = 9$ and the coefficient of correlation is $\rho_{XY} = \frac{1}{2}$.
 Calculate $E(Z)$ where $Z = (X - Y)^2$. [4]

- B5.** (a) A random vector (X, Y) is normally distributed with parameters (μ_X, μ_Y) ; σ_X ; σ_Y and ρ_{XY} .

- (i) Calculate the conditional density function of Y given X, $f_{Y|X}(x, y)$. [8]
 (ii) Hence or otherwise, calculate the conditional expectation $E[Y|X = x]$. [3]

- (b) The random vector (X, Y) has the following characteristics:

$$E[X] = -2; E[Y] = 3 \text{ and the covariance matrix is } K = \begin{pmatrix} 16 & 12 \\ 12 & 25 \end{pmatrix}$$

Calculate the joint density function $f_{X,Y}(x, y)$. [6]

- (c) The joint probability density function of a random vector (X, Y) is given by

$$f_{X,Y}(x, y) = \frac{1}{\pi} e^{-\frac{1}{2}(x^2 + 2xy + 5y^2)}$$

Calculate $f_X(x)$ and ρ_{XY} . [5,3]

- B6.** (a) A random variable X is normally distributed with mean μ and variance σ^2 .

Calculate the characteristic function of X. [6]

Hence or otherwise calculate $E[X]$ and $\text{Var}[X]$ if they exist. [3,3]

- (b) Given that the characteristic function of a continuous random variable X is

$$\Theta_X(t) = \frac{e^{itm}}{1 + \frac{\sigma^2 t^2}{2}}$$

Find the density function of X and calculate $E(X)$ and $\text{Var}(X)$ as functions of m and σ . [7,3,3]

- B7.** (a) A sequence X_1, X_2, \dots , of random variables satisfies $E[X_k^2] < \alpha^2, \forall k = 1, 2, \dots$
Given also that X_k and X_{k+1} are independent
- (i) Find the upper bound of the probability

$$P\left(\left|\frac{X_k + X_{k+1}}{2} - \frac{E(X_k) + E(X_{k+1})}{2}\right| \geq \sigma_k + \sigma_{k+1}\right)$$
where $\sigma_j = \sqrt{\text{Var}(X_j)}, \forall j = 1, 2, \dots$ [7]
- (ii) Prove that the sequence satisfies Markov's law of large numbers. [6]
- (b) A random variable X has a Poisson distribution with parameter λ . Prove that when $\lambda \rightarrow \infty$, then

$$Z = \frac{X - \lambda}{\sqrt{\lambda}}$$
 has a standard normal distribution. [6]
- (c) Let f be an increasing function and assume that $E[f(x)]$ exists, where X is a nonnegative random variable.
Prove that $\forall \varepsilon, P(X \geq \varepsilon) \leq \frac{E[f(x)]}{f(\varepsilon)}$. [6]
- B8.** (a) Calculate the moment generating function of a random variable X whose probability density function is

$$f_X(y) = \frac{1}{b\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{y-\mu}{b}\right)^2}, \quad -\infty < y < \infty.$$
 [7]
- (b) Calculate $E(X)$ and $\text{Var}(X)$ if X is a random variable with density

$$f_X(x) = \begin{cases} \theta e^{-\theta x}, & x > 0 \\ 0 & \text{otherwise} \end{cases}$$
 [3,4]
- (c) A random vector (X, Y) has joint density function

$$f_{X,Y}(x, y) = \begin{cases} e^{-(x+y)}, & \text{if } x, y \geq 0 \\ 0 & \text{otherwise} \end{cases}$$
- (i) Calculate $E(XY)$. [6]
- (ii) Determine whether or not X and Y are independent. [5]

END OF QUESTION PAPER