

NATIONAL UNIVERSITY OF SCIENCE AND TECHNOLOGY

APPLIED PHYSICS DEPARTMENT

SPH 1105 – ELECTRICITY AND MAGNETISM

SUPPLEMENTARY EXAMINATION

BSC HONOURS PART I

AUGUST 2004

DURATION: 3 HOURS

ANSWER ALL PARTS OF QUESTION 1-IN SECTION A AND ANY THREE QUESTIONS FROM SECTION B. SECTION A CARRIES 40 MARKS AND SECTION B CARRIES 60 MARKS .

Gravitational constant $G$	$= 6.67 \times 10^{-11} \text{ Nm}^2 \text{ kg}^{-2}$
Permittivity of free space $\epsilon_0$	$= 8.85 \times 10^{-12} \text{ Fm}^{-1}$
Permeability of free space $\mu_0$	$= 1.26 \times 10^{-6} \text{ Hm}^{-1}$
Electron mass $m_e$	$= 9.11 \times 10^{-31} \text{ kg}$
Proton mass $m_p$	$= 1.66 \times 10^{-27} \text{ kg}$
Charge on an electron $e$	$= 1.60 \times 10^{-19} \text{ C}$

SECTION A

1. (a) Equal electric charges of  $3 \mu\text{C}$  each are placed at the vertices of an equilateral triangle whose sides are  $2.5 \text{ cm}$  in length Calculate the electric field at each vertex of the triangle. [4]
- (b) A dielectric material is placed between the plates of a charged capacitor. Explain the effect on
  - (i) electric field between the plates
  - (ii) capacitance [4]
- (c) Two point charges  $q_1 = 2.5 \mu\text{C}$  and  $q_2 = -2.5 \mu\text{C}$  are separated by  $5 \text{ mm}$ . What is the dipole moment of these two charges? Sketch the pair and indicate the direction of the dipole moment. [6]
- (d) A galvanometer of internal resistance  $20\Omega$  reads full scale deflection when  $45 \text{ mA}$  passes through it. Design an ammeter to read up to  $2.0 \text{ A}$  [4]
- (e) (i) What is a solenoid? [2]
  - (ii) Find the magnetic flux at the centre of a long, tightly wound solenoid of length  $50 \text{ cm}$  and radius  $5 \text{ cm}$  carrying a current of  $300 \text{ mA}$ . The number of turns is  $300$ . [4]

- (f) Derive the balance condition of a Wheatstone Bridge. Explain how it can be used to calculate an unknown resistance. [4]
- (g) Explain briefly how static electricity can be used in Xerography. [4]
- (h) Find the magnetic field induction at the centre of a square current loop of side 2m carrying a current of 2 A as shown in Figure 1 below. [4]




Figure 1

- (i) A long wire parallel to the x-axis carries a current of 8 A in the direction of increasing x. There is a uniform magnetic field of magnitude 1.8 T in the positive y- direction. Find the force per unit length on the wire. [4]

### SECTION B

2. (a) Distinguish between reactance and impedance. Illustrate your answer with relevant expressions. [5]
- (b) In a certain LCR circuit  $X_C = 16 \Omega$ ,  $X_L = 4 \Omega$  at some frequency  $\omega$ . The resonance frequency  $\omega_0 = 10^4$  rad/s.  
 (i) Find L and C. [4]
- If  $R = 5 \Omega$  and  $\varepsilon_{\max} = 26 \text{ V}$ , find  
 (ii) the Q value, [3]  
 (iii) the maximum current. [3]
- (c) Compute by direct integration the area under the curve  $\sin^2 \omega t$  from  $t = 0$  to  $t = T = 2\pi/\omega$  and show that it is equal to  $\frac{1}{2}T$ . [5]
3. (a) Define  
 (i) the ampere and  
 (ii) the coulomb. [4]

- (b) Describe the motion of a positive charge if it is:
- released into a uniform magnetic field of intensity  $B$  at an angle  $\theta$  to the field. [4]
  - released into the same field with a velocity directly perpendicular to the field. [4]

- (c) A current-carrying wire induces a magnetic field into the surrounding space, given by Biot – Savart Law:

$$dB = \frac{\mu_0 I dx \hat{r}}{\pi r^2}$$

Explain all terms in the above equation [6]

- (d) Use this law to derive the magnitude of the magnetic field due a thin wire of infinite length carrying a current  $I$  [6]

4. (a) State Kirchoff's laws. [2]

- (b) In Figure 2 below, find
- the current in each resistor [3]
  - the potential difference between  $a$  and  $b$  [3]
  - the power dissipated in each resistor. [3]
- Take  $E_1 = 6.0 \text{ V}$ ,  $E_2 = 5.0 \text{ V}$ ,  $E_3 = 4.0 \text{ V}$ ,  $R_1 = 100 \Omega$  and  $R_2 = 50 \Omega$




Figure 2

- (c) A  $1.0\mu\text{f}$  capacitor with an initial stored energy of  $0.50\text{J}$  is discharged through a  $1.0\text{M}\Omega$  resistor.
- What is the initial charge on the capacitor. [3]
  - What is the current through the resistor when the discharge starts. [3]
  - Determine  $V_C$ , the voltage across the capacitor and  $V_R$ , the voltage across the resistor as a function of time. [3]
5. (a) (i) State Gauss's law and explain the physical meaning of all quantities involved. [4]

(ii) Use this law to derive an expression for the electric field due to a spherical distribution of charges. [4]

(b) Figure 3 shows a charge  $+q$  arranged as a uniform non-conducting sphere of radius  $a$  and placed at the centre of a spherical conducting shell of wider radius  $b$  and outer radius  $c$ . The outer shell carries a charge of  $-q$ . Find  $E(r)$




Figure 3

- (i) within the sphere ( $r < a$ ) [4]
- (ii) between the sphere and the shell ( $a < r < b$ ) [2]
- (iii) inside the shell ( $b < r < c$ ) [2]
- (iv) outside the shell ( $r > c$ ) [2]
- (v) what charges appear on the inner and outer surfaces of the shell. [2]

6. (a) Distinguish between potential and potential difference. [4]
- (b) What is an equi-potential surface? [4]
- (c) Three positive  $2 \mu\text{C}$  point charges are at the corners of a square of side  $3\text{m}$ . [3]
- (i) What is the potential  $V$  at the fourth unoccupied corner of the square. [3]
  - (ii) How much work is needed to bring up a fourth positive charge of  $2\mu\text{C}$  and place it at the fourth corner of the square? [3]
- (d) Find the potential on the axis of a disk of uniform surface charge density  $\sigma$ . [6]

END OF PAPER